## Emergency Medical Technician Psychomotor Examination CARDIAC ARREST MANAGEMENT / AED

Candidate:	Examiner:	
Date:	Start Time:	End Time:

Takes or verbalizes appropriate body substance isolation precautions		
Determines the scene situations is safe	1	
Attempts to question bystanders about arrest events		
Checks patient responsiesness	1	
NOTE: The examiner must not inform the candidate, "The patient is unresponsive		
Assessess patient for signs of breathing (observes the patient and determines the absence of	1	
breathing or abnormal breathing (gasping and agonal respirations)		
NOTE: The examiner must not inform the candidate, "The patient is apneic," or "The patient		
has gasping, agonal, respriations."		
Checks carotid pulse [no more than 10 seconds]	1	
Note:The examiner must not inform the candidate, "The patient is pulseless."		
Immedicately begins chest compressions [adequate deth and rate;	1	
allows the chest to recoil completeley]		
Requests additional EMS response	1	
forms 2 minutes of high quality, 1-rescuer adult CPR		
Adequate depth and rate	5	
Correct compression-to-ventilation ratio		
Allows the chest to recoil completely		
Adequate volumes for each breath		
Minimal interruptions of less than 10 seconds throughout	7	
NOTE: After 2 minutes (5 cycles), patient is assessed and second rescuer operates AED.		
resumes compressions while candidate		
Turns-on power to AED	1	
Follows prompts and correctly attaches AED to patient	1	
Stops CPR and ensures all individuals are clear of the patient during rhythm analysis	1	
Ensures that all individuals are clear of the patient and delivers shock from AED		
Immediately directs rescuer to resume chest compressions	1	
Total	18	

You must factually document your rationale for checking any of the above critical items on this form (below or turn sheet over).

Failure to take or verbalize appropriate body substance isolation precautions	
Failure to immediately begin chest compressions as soon as pulselessness is confir	med
Failure to deliver shock in a timely manner	
Interrupts CPR for more than 10 seconds at any point	
Failure to demonstrate acceptable high-quality, 1-rescuer adult CPR	
Failure to operate the AED properly	
Failure to correctly attach the AED to the patient	
Failure to assure that all individuals are clear of patient during rhythm analysis obs	serves]
and [verbalizes "All clear" and before delivering shock(s)	-
Failure to immediately resume compressions after shock delivered	
Failure to manage the patient as a competent EMT	
Exhibits unacceptable affect with patient or other personnel	
Uses or orders a dangerous or inappropriate intervention	
oses of orders a dangerous of mappropriate intervention	